

# Cyclodextrins and Their Complexes

Chemistry, Analytical Methods, Applications

*Edited by Helena Dodziuk*



WILEY-  
VCH

WILEY-VCH Verlag GmbH & Co. KGaA



**Cyclodextrins and Their  
Complexes**

*Edited by  
Helena Dodziuk*

## ***Related Titles***

Elias, H.-G.

### **Macromolecules**

**Volume 1: Chemical Structures and Syntheses**

2005

ISBN 3-527-31172-6

Munk, P., Aminabhavi, T. M.

### **Introduction to Macromolecular Science**

2002

ISBN 0-471-41716-5

Bladon, C.

### **Pharmaceutical Chemistry**

**Therapeutic Aspects of Biomacromolecules**

2002

ISBN 0-471-49636-7

# **Cyclodextrins and Their Complexes**

Chemistry, Analytical Methods, Applications

*Edited by Helena Dodziuk*



**WILEY-  
VCH**

WILEY-VCH Verlag GmbH & Co. KGaA

#### The Editor

**Prof. Dr. Helena Dodziuk**

Inst. of Physical Chemistry  
Polish Academy of Sciences  
Kasprzaka 44  
01-224 Warsaw  
Poland  
and

Graduate College ICM-IWR  
Warsaw (Poland) – Heidelberg (Germany)

■ All books published by Wiley-VCH are carefully produced. Nevertheless, authors, editors, and publisher do not warrant the information contained in these books, including this book, to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

**Library of Congress Card No.:** applied for  
**British Library Cataloguing-in-Publication Data:**  
A catalogue record for this book is available from the British Library.

**Bibliographic information published by  
Die Deutsche Bibliothek**  
Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data is available in the Internet at <http://dnb.ddb.de>.

© 2006 WILEY-VCH Verlag GmbH & Co.  
KGaA, Weinheim

All rights reserved (including those of translation into other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Printed in the Federal Republic of Germany.  
Printed on acid-free paper.

**Typesetting** Asco Typesetters, Hong Kong  
**Printing** betz-druck GmbH, Darmstadt  
**Binding** Schäffer GmbH, Grünstadt  
**Cover Design** Grafik-Design Schulz,  
Fußgönheim

**ISBN-13:** 978-3-527-31280-1

**ISBN-10:** 3-527-31280-3

## Contents

**Preface** XIII

**List of Contributors** XV

<b>1</b>	<b>Molecules with Holes – Cyclodextrins</b>	<b>1</b>
	<i>Helena Dodziuk</i>	
1.1	Introduction	1
1.2	Cyclodextrin Properties	17
1.3	Cyclodextrin Nonrigidity	20
1.4	Models of Chiral Recognition by Cyclodextrins	21
	References	26
<b>2</b>	<b>Modification Reactions of Cyclodextrins and the Chemistry of Modified Cyclodextrins</b>	<b>31</b>
	<i>Kenjiro Hattori and Hiroshi Ikeda</i>	
2.1	Scope of This Chapter	31
2.2	Modification Reactions of Cyclodextrins	31
2.2.1	Modification Reactions at the Primary Side	32
2.2.1.1	Mono-modification at the C6-Position	32
2.2.1.2	Per-modification at the C6-Position	35
2.2.1.3	Multi-modification at the C6-Position	36
2.2.2	Modification Reactions at the Secondary Side	38
2.2.2.1	Mono-modification at the C2-Position	38
2.2.2.2	Mono-modification at Any One of the C2-, C3-, or C6-Positions	38
2.2.2.3	Per-modification at the C3-, C2-, or C6-Position	39
2.2.3	Per-modification Reactions at All Three Positions	40
2.2.4	Enzymatic Modification Reactions of Cyclodextrins	40
2.2.5	Construction Reactions for Cyclodextrin Ring Formation	41
2.3	Chemistry of Modified Cyclodextrins	41
2.3.1	Cyclodextrin Dimers and Trimers	41
2.3.2	Charged Cyclodextrins	45
2.3.3	Chemosensors Using Modified Cyclodextrins	47
2.3.4	Cyclodextrin Analogues	50

2.3.5	Cyclodextrins Conjugated with Other Kinds of Hosts	51
2.3.6	Cyclodextrin–Peptide Conjugates	53
2.3.7	Cyclodextrin–Saccharide Conjugates	55
2.3.8	Metallo-cavitands Using Modified Cyclodextrins	57
	References	59
<b>3</b>	<b>Polymers Involving Cyclodextrin Moieties</b>	<b>65</b>
	<i>Akira Harada, Akihito Hashidzume, and Masahiko Miyauchi</i>	
3.1	Supramolecular Polymers Formed by Cyclodextrin Derivatives	65
3.1.1	Introduction	65
3.1.2	Preparation of Mono-substituted Cyclodextrins	66
3.1.3	Formation of Inclusion Complexes	68
3.1.4	Polymer Formation by Intermolecular Interactions	69
3.1.5	Supramolecular Dimers	70
3.1.6	Formation of Supramolecular Trimers	71
3.1.7	Cyclic Daisy Chain	72
3.1.8	Supramolecular Polymers	74
3.1.9	Poly[2]rotaxanes (Daisy Chain)	76
3.1.10	Helical Supramolecular Polymers	77
3.1.11	Alternating $\alpha$ -, $\beta$ -Cyclodextrin Supramolecular Polymers	82
3.1.12	Supramolecular [2]rotaxane Polymer	85
3.1.13	Conclusion	86
3.2	Supramolecular Complexes of Polymers Bearing Cyclodextrin Moieties with Guest Molecules	89
	References	89
<b>4</b>	<b>Cyclodextrin Catalysis</b>	<b>93</b>
	<i>Makoto Komiyama and Eric Monflier</i>	
4.1	Introduction	93
4.2	Covalent Catalysis	94
4.3	General Acid–Base Catalysis by OH Groups	95
4.4	Noncovalent Catalysis	96
4.4.1	Regulation of the Mutual Conformation of Reactants	97
4.4.2	Regulation of Photoreactions	99
4.4.3	Use of the CyD Cavity as a Specific Reaction Field	99
4.5	Catalysis by Chemically Modified CyD	101
4.6	Phase-transfer Catalysis	102
4.7	Conclusion	103
	References	104
<b>5</b>	<b>Chromatographic Studies of Molecular and Chiral Recognition</b>	<b>106</b>
	<i>Monika Asztemborska and Anna Bielejewska</i>	
5.1	Introduction	106
5.2	Determination of the Stoichiometry and Stability of the Complexes	108
5.3	Thermodynamics	116
	References	117



<b>6</b>	<b>The Application of Cyclodextrins for Enantioseparations</b>	<b>119</b>
	<i>Bezhan Chankvetadze</i>	
6.1	Introduction	119
6.2	Gas Chromatography	120
6.3	High-performance Liquid Chromatography and Related Techniques	123
6.4	Super (Sub)-critical Fluid Chromatography	124
6.5	Capillary Electrophoresis	125
6.6	Capillary Electrochromatography	141
6.7	Lab-on-a-chip Technologies	143
6.8	Future Trends	144
	References	144
<b>7</b>	<b>Crystallographic Study of Cyclodextrins and Their Inclusion Complexes</b>	<b>147</b>
	<i>Kazuaki Harata</i>	
7.1	Introduction	147
7.2	Native CyDs	147
7.2.1	Structure of the Glucose Unit	148
7.2.2	Average Structural Characteristics of the Macrocyclic Ring	150
7.2.3	Hydrogen Bonds	153
7.3	Modified CyDs	153
7.3.1	Methylated CyDs	153
7.3.2	Acylated CyDs	155
7.3.3	Monosubstituted CyDs	156
7.3.4	Disubstituted CyDs	160
7.3.5	Capped CyDs	160
7.3.6	CyDs with Nonglucose Units	162
7.4	Crystalline Inclusion Complexes	163
7.4.1	Host–Guest Interactions	163
7.4.2	$\alpha$ -CyD Complexes	166
7.4.3	$\beta$ -CyD Complexes	170
7.4.4	Complexes of Larger CyDs	176
7.4.5	Complexes of Methylated CyDs	177
7.4.6	Complexes with Metals and Organometals	182
7.5	Molecular Recognition in the Solid State	186
7.5.1	Selectivity for Geometrical Isomers (see also Chapter 5)	186
7.5.2	Chiral Recognition (see also Chapters 1, 5, 6)	186
7.6	Concluding Remarks	191
	References	191
<b>8</b>	<b>Microcalorimetry</b>	<b>199</b>
	<i>Mikhail V. Rekharsky and Yoshihisa Inoue</i>	
8.1	Introduction	199
8.2	Application of Differential Scanning Calorimetry in Cyclodextrin and Drug Studies	200
8.3	Application of Isothermal Titration Calorimetry in Cyclodextrin and Drug Studies	202

- 8.4 Determination of Thermodynamic Parameters for Stoichiometric 1:1 Complexation by Cyclodextrin 206
- 8.5 Determination of Thermodynamic Parameters for 1:2 Guest–Host Complexation of Cyclodextrin 212
- 8.6 A Thermodynamic View for Globally Understanding Molecular Recognition 215
- 8.7 Enthalpy–Entropy Compensation as a Practical Tool for a Global Understanding of Molecular Recognition 216
- 8.8 Grunwald Theory on Enthalpy–Entropy Compensation 219
- 8.9 Isotope Effect: Implications for the Comparison of Complexation Thermodynamics in H<sub>2</sub>O versus D<sub>2</sub>O 219
- 8.10 A Theoretical Approach to Enthalpy–Entropy Compensation in Molecular Recognition 222
- 8.11 Conclusions 222  
References 223
- 9 NMR of Cyclodextrins and Their Complexes 231**  
*Andrzej Ejchart and Wiktor Koźmiński*
- 9.1 NMR Methodology 231
- 9.1.1 Standard NMR Spectra 231
- 9.1.2 Nuclear Magnetic Relaxation 234
- 9.1.3 Dipolar Interactions and NOE 235
- 9.1.4 Molecular Dynamics 236
- 9.1.5 2D Spectra 237
- 9.1.6 Translational Self-diffusion 241
- 9.2 Application of NMR to Cyclodextrins and Their Complexes 242
- 9.2.1 NMR Parameters and the Complexing Ability of Cyclodextrins and Their Derivatives 242
- 9.2.2 Stoichiometry 243
- 9.2.3 Determination of Association Constants 244
- 9.2.4 Host–Guest Orientation and Geometry of CyD Inclusion Complexes 246
- 9.2.5 Chiral Recognition 248
- 9.2.6 NMR Experiments in the Solid State 250  
References 252
- 10 Other Physicochemical Methods 255**
- 10.1 Introductory Remarks 255  
*Helena Dodziuk*  
References 256
- 10.2 Mass Spectrometry of CyDs and Their Complexes 257  
*Witold Danikiewicz*
- 10.2.1 Introduction – Mass Spectrometry Techniques Used for Studying Large Molecules and Their Complexes 257
- 10.2.2 Mass Spectrometry of Free CyDs 259

10.2.2.1	Fast Atom Bombardment (FAB) and Liquid-matrix Secondary-ion Mass Spectrometry (LSIMS)	259
10.2.2.2	Electrospray Ionization (ESI)	260
10.2.2.3	Matrix-Assisted Laser Desorption Ionization (MALDI)	262
10.2.3	Mass Spectrometry of Inclusion Complexes of CyDs	265
10.2.3.1	Fast Atom Bombardment (FAB) and Liquid-matrix Secondary Ion Mass Spectrometry (LSIMS)	267
10.2.3.2	Electrospray Ionization (ESI)	267
10.2.3.3	Matrix-assisted Laser Desorption Ionization (MALDI)	273
	References	273
10.3	Studies of Cyclodextrin Inclusion Complexes by Electronic (UV-Vis Absorption and Emission) Spectroscopy	276
	<i>Gottfried Grabner</i>	
10.3.1	Introduction	276
10.3.2	Steady-state Spectroscopies	277
10.3.2.1	UV-Vis Absorption Spectroscopy	277
10.3.2.2	Fluorescence Emission Spectroscopy	279
10.3.2.3	Room-temperature Phosphorescence (RTP)	281
10.3.3	Time-resolved Spectroscopies	282
10.3.3.1	Excited-state Absorption	282
10.3.3.2	Complexation Kinetics	283
10.3.3.3	Host Structural Flexibility	285
10.3.3.4	Ultrafast Studies of Guest Dynamics	286
	References	286
10.4	Circular Dichroism of Cyclodextrin Complexes	289
	<i>Daniel Krois and Udo H. Brinker</i>	
10.4.1	General Introduction to Circular Dichroism	289
10.4.2	Types of CD Observed for Cyclodextrin Complexes	289
10.4.3	Application of CD Spectroscopy for the Characterization of Cyclodextrin Complexes	292
10.4.3.1	Determination of Complex Stability and Stoichiometry by CD	292
10.4.3.2	Determination of Complex Geometry by CD	295
10.4.4	Conclusions	296
	References	296
10.5	Electrochemistry of Cyclodextrins	298
	<i>Renata Bilewicz and Kazimierz Chmurski</i>	
10.5.1	Introduction	298
10.5.2	Electrochemistry in Homogeneous Solutions	304
10.5.3	Electrochemistry of Cyclodextrin Thin Films	304
10.5.4	Summary and Prospects	315
	References	316
10.6	Visualization of Cyclodextrins in Supramolecular Structures by Scanning Probe Microscopy	319
	<i>Masashi Kunitake and Akihiro Ohira</i>	

- 10.6.1 Introduction 319
- 10.6.2 STM or AFM Observations of a CyD Nanotube in Air 320
- 10.6.3 STM Observation of CyDs in Aqueous Solution 324
- 10.6.4 Dynamic Force Spectroscopy 328
- 10.6.5 Conclusion 329
- References 330

**11 Modeling of CyDs and Their Complexes 333**

*Helena Dodziuk*

- 11.1 Introduction 333
- 11.2 Quantum Chemical Calculations 337
- 11.3 Molecular Mechanics Cyclodextrin Studies 341
- 11.4 Dynamic Simulations CyD Studies 346
- 11.5 Conclusions 351
- References 352

**12 Rotaxane and Catenane Structures Involving Cyclodextrins 356**

*Konstantina Yannakopoulou and Irene M. Mavridis*

- 12.1 Introduction 356
- 12.2 Pseudorotaxanes 357
- 12.3 Rotaxanes 360
- 12.4 Applications 362
- 12.5 Catenanes 365
- References 367

**13 Large-ring Cyclodextrins 370**

*Haruhisa Ueda and Tomohiro Endo*

- 13.1 Introduction 370
- 13.2 Production of LR-CyDs 371
- 13.3 Isolation and Purification Procedures 372
- 13.4 Physicochemical Properties and Structures of LR-CyDs 373
- 13.5 Inclusion Complex Formation 375
- 13.6 Applications of LR-CyDs 378
- References 378

**14 Pharmaceutical Applications of Cyclodextrins and Their Derivatives 381**

*Kaneto Uekama, Fumitoshi Hirayama, and Hidetoshi Arima*

- 14.1 Introduction 381
- 14.2 Some Characteristics of CyDs as Drug Carriers 383
  - 14.2.1 Hydrophilic CyDs 384
  - 14.2.2 Hydrophobic CyDs 387
  - 14.2.3 Ionizable CyDs 389
- 14.3 Improvements in the Pharmaceutical Properties of Drugs 390
  - 14.3.1 Bioavailability 390
  - 14.3.2 Chemical and Physical Stability 393
    - 14.3.2.1 In Solution 393

14.3.2.2	In the Solid State	396
14.3.3	Release Control	398
14.3.3.1	Immediate-release	398
14.3.3.2	Prolonged-release	399
14.3.3.3	Modified-release	400
14.3.3.4	Delayed-release	401
14.3.4	Enhancement of CyD Action by Combined Use with Additives	402
14.4	Potential Use of CyDs in Peptide and Protein Formulation	405
14.4.1	Absorption Enhancement by Hydrophilic CyDs	405
14.4.2	Sustained-release of Peptide and Protein Drugs	407
14.4.3	Inhibitory Effects of CyDs on Aggregate Formation of Polypeptides	409
14.5	Site-specific Drug Delivery	410
14.5.1	Colonic Delivery	411
14.5.2	Application in Gene Therapies	415
14.6	Perspectives	417
	References	418
<b>15</b>	<b>Cyclodextrins in Dispersed Systems</b>	<b>423</b>
	<i>Laury Trichard, Dominique Duchêne, and Amélie Bochot</i>	
15.1	Introduction	423
15.2	Dispersed Systems Involving CyDs	432
15.2.1	Emulsions	432
15.2.2	Microparticles	433
15.2.2.1	Microcapsules	433
15.2.2.2	Microspheres	435
15.2.3	Nanoparticles	436
15.2.3.1	Polymeric Nanoparticles	436
15.2.3.2	Self-assembled Polymeric Nanoparticles	436
15.2.3.3	Amphiphilic Cyclodextrin Nanoparticles	437
15.2.4	Lipidic Vesicles	438
15.2.4.1	Liposomes and Niosomes Prepared With Native or Water-soluble Modified CyDs	438
15.2.4.2	Lipidic Vesicles and Amphiphilic CyDs	440
15.3	The Potentialities of CyDs in Dispersed Systems	440
15.3.1	Conventional Uses of CyDs in Dispersed Systems	441
15.3.1.1	Enhancement of Drug Solubility and Loading	441
15.3.1.2	Enhancement of Drug Stability	441
15.3.2	Cyclodextrins as New Materials for Particle Preparation	441
15.3.3	CyDs to Control Drug Release Profile	442
15.3.3.1	Microparticles	442
15.3.3.2	Nanoparticles	442
15.3.3.3	Liposomes	443
15.3.4	Toxicity and Efficacy Evaluation	444
15.3.4.1	Microparticles	444
15.3.4.2	Polymeric Nanoparticles	444

15.3.4.3	Amphiphilic CyD Nanoparticles	444
15.3.4.4	Lipidic Vesicles	445
15.4	Conclusions	445
	References	446
<b>16</b>	<b>Applications Other Than in the Pharmaceutical Industry</b>	<b>450</b>
16.1	Introductory Remarks	450
	<i>Helena Dodziuk</i>	
	References	451
16.2	CyD Applications in Food, Cosmetic, Toiletry, Textile and Wrapping Material Fields	452
	<i>Hitoshi Hashimoto</i>	
16.2.1	In Foods	452
16.2.1.1	Stabilization by Powdering	452
16.2.1.2	Taste Modification	453
16.2.1.3	Anti-oxidation, Stabilization, and Improvement of Bioavailability	453
16.2.1.4	Solubilization of Insoluble Substances	455
16.2.2	In Cosmetics and Toiletries	455
16.2.3	In Textiles	456
16.2.4	In Wrapping Materials	457
16.2.5	Future Prospects	457
	References	458
16.3	Application of Cyclodextrins in Agrochemistry	459
	<i>Esmeralda Morillo</i>	
	Conclusions	465
	References	466
16.4	Cyclodextrins in Molecular Devices	468
	<i>Renata Bilewicz and Kazimierz Chmurski</i>	
16.4.1	Introduction	468
16.4.2	Electrochemically Controlled Molecular Machines	469
16.4.3	Applications of CyDs in Bioelectrocatalysis	470
	References	472
	<b>Index</b>	<b>475</b>

## Preface

Cyclodextrins and their complexes constitute a fascinating subject that, due to its complexity, lacked an extended coverage in a monograph for long time. We are happy to bridge this gap with this book. The field has grown immensely, thus discussing or even mentioning many valuable works was not possible in the limited volume. However, we hope to have given an extensive description of all important aspects of research and applications involving cyclodextrins.

The creation of this book would not be possible without the assistance of several people deserving acknowledgement. They have helped by précising its content, choosing contributors, reviewing chapters, and in other ways. I am especially indebted to Prof. H. Ueda, Prof. R. Bilewicz and Prof. K. Harata for their involvement in the project. The critical remarks on specific chapters by Prof. A. Teme-riusz, Prof. M. Kańska, Prof. J. Frelek, Prof. A. Holas, Prof. M. Jaszuński, Dr. R. Nowakowski and Dr. K. Chmurski are gratefully acknowledged. I would like also to thank Wacker, GmbH for a generous gift of cyclodextrins that helped me to carry out cyclodextrin experimental studies, and Oxford University that has enabled my literature research at Kebble College, leading to my deeper understanding of supramolecular complexes.

Warsaw, January 2006

*Helena Dodziuk*





## List of Contributors

### **Hidetoshi Arima**

Graduate School of Pharmaceutical Sciences  
Kumamoto University  
5-1, Oe-honmachi  
Kumamoto 862-0973  
Japan

### **Monika Asztemborska**

Institute of Physical Chemistry  
Polish Academy of Sciences  
01-224 Warsaw  
Kasprzaka 44/52  
Poland

### **Anna Bielejewska**

Institute of Physical Chemistry  
Polish Academy of Sciences  
Kasprzaka 44/52  
01-224 Warsaw  
Poland

### **Renata Bilewicz**

Department of Chemistry  
University of Warsaw  
Pasteura 1  
02-093 Warsaw  
Poland

### **Amélie Bochot**

Laboratoire de Pharmacie galénique  
UMR CNRS 8612  
Centre d'Etudes Pharmaceutiques  
Université Paris Sud  
5 rue J.B. Clément  
F-92290 Châtenay-Malabry  
France

### **Udo H. Brinker**

Institut für Organische Chemie der  
Universität Wien  
Währinger Straße 38  
1090 Wien  
Austria

### **Bezhan Chankvetadze**

Molecular Recognition and Separation Science  
Laboratory  
School of Chemistry  
Tbilisi State University, Chavchavadze Ave 1  
380028 Tbilisi  
Georgia

### **Kazimierz Chmurski**

Department of Chemistry  
University of Warsaw  
Pasteura 1  
02-093 Warsaw  
Poland

### **Witold Danikiewicz**

Institute of Organic Chemistry  
Polish Academy of Sciences  
01-224 Warsaw  
Kasprzaka 44/52  
Poland

### **Helena Dodziuk**

Institute of Physical Chemistry  
Polish Academy of Sciences  
Kasprzaka 44/52  
01-224 Warsaw  
Poland  
and  
Graduate College ICM-IWR  
Warsaw – Heidelberg

### **Dominique Duchêne**

Laboratoire de Pharmacie galénique  
UMR CNRS 8612  
Centre d'Etudes Pharmaceutiques  
Université Paris Sud  
5 rue J.B. Clément  
F-92290 Châtenay-Malabry

**Andrzej Ejchart**

Institute of Biochemistry and Biophysics  
Polish Academy of Sciences  
Pawińskiego 5A  
02-106 Warszawa  
Poland

**Tomohiro Endo**

Department of Physical Chemistry  
Hoshi University  
4-41, Ebara 2-chome  
Shinagawa-ku  
Tokyo 142-8501  
Japan

**Gottfried Grabner**

Max F. Perutz Laboratories  
Department of Chemistry  
University of Vienna  
Campus Vienna Biocenter 5  
1030 Vienna  
Austria

**Akira Harada**

Laboratory of Supramolecular Science  
Department of Macromolecular Science  
Graduate School of Science  
Osaka University  
Toyonaka  
Osaka 560-0043  
Japan

**Kazuaki Harata**

Biological Information Research Center  
National Institute of Advanced Industrial  
Science and Technology  
Central 6, 1-1-1 Higashi  
Tsukuba  
Ibaraki 305-8566  
Japan

**Akihito Hashidzume**

Laboratory of Supramolecular Science  
Department of Macromolecular Science  
Graduate School of Science  
Osaka University  
Toyonaka  
Osaka 560-0043  
Japan

**Hitoshi Hashimoto**

Bio Research Corporation of Yokohama  
Yokohama Kanazawa High-Tech Center  
Techno-Core  
1-1-1 Fukuura  
Kanazawa-Ku  
Yokohama 236-0004  
Japan

**Kenjiro Hattori**

Department of Nanochemistry,  
Faculty of Engineering  
Tokyo Polytechnic University  
1583 Iiyama  
Atsugi 243-0297  
Japan

**Fumitoshi Hirayama**

Graduate School of Pharmaceutical Sciences  
Kumamoto University  
5-1, Oe-honmachi  
Kumamoto 862-0973  
Japan

**Hiroshi Ikeda**

Department of Bioengineering,  
Graduate School of Bioscience and  
Biotechnology  
Tokyo Institute of Technology  
4259-B-44 Nagatsuta-cho, Midori-ku  
Yokohama 226-8501  
Japan

**Yoshihisa Inoue**

Department of Applied Chemistry  
Osaka University  
2-1 Yamada-oka  
Suita 565-0871  
Japan

**Makoto Komiyama**

Research Center for Advanced Science and  
Technology  
The University of Tokyo  
4-6-1 Komaba  
Meguro-ku  
Tokyo 153-8904  
Japan

**Wiktor Koźmiński**

Department of Chemistry  
Warsaw University  
Pasteura 1  
02-093 Warszawa  
Poland

**Daniel Krois**

Institut für Organische Chemie der  
Universität Wien  
Währinger Straße 38  
1090 Wien  
Austria

**Masashi Kunitake**

Graduate School of Science and Technology  
Department of Applied Chemistry and  
Biochemistry  
Kumamoto University  
39-1 Kurokami 2-chome  
Kumamoto 860-8555  
Japan

**Irene M. Mavridis**

Institute of Physical Chemistry  
N.C.S.R. "Demokritos"  
P.O. Box 60228  
153 10 Aghia Paraskevi  
Athens  
Greece

**Masahiko Miyauchi**

Laboratory of Supramolecular Science  
Department of Macromolecular Science  
Graduate School of Science  
Osaka University  
Toyonaka  
Osaka 560-0043  
Japan

**Eric Monflier**

Faculté des Sciences J. Perrin, Université  
d'Artois  
Rue Jean Souvraz/Sac postal 18  
F-62307 Lens Cédex

**Esmeralda Morillo**

Instituto de Recursos Naturales y Agrobiología  
de Sevilla  
Avda. Reina Mercedes 10  
Apdo 1052  
41080-Sevilla  
Spain

**Akihiro Ohira**

Graduate School of Materials Science  
Nara Institute of Science and Technology  
8916-5, Takayama, Ikoma  
Nara 630-0101  
Japan

**Mikhail V. Rekharsky**

Entropy Control Project  
ICORP  
Japan Science and Technology Agency  
4-6-3 Kamishinden  
Toyonaka 565-0085  
Japan

**Laury Trichard**

Laboratoire de Pharmacie galénique  
UMR CNRS 8612  
Centre d'Etudes Pharmaceutiques  
Université Paris Sud  
5 rue J.B. Clément  
F-92290 Châtenay-Malabry

**Hauhisa Ueda**

Department of Physical Chemistry  
Hoshi University  
4-41, Ebara 2-chome  
Shinagawa-ku  
Tokyo 142-8501  
Japan

**Kaneto Uekama**

Graduate School of Pharmaceutical Sciences  
Kumamoto University  
5-1, Oe-honmachi  
Kumamoto 862-0973  
Japan

**Konstantina Yannakopoulou**

Institute of Physical Chemistry  
N.C.S.R. "Demokritos"  
P.O. Box 60228  
153 10 Aghia Paraskevi  
Athens  
Greece



# 1


## Molecules with Holes – Cyclodextrins

Helena Dodziuk

### 1.1 Introduction


Cyclodextrins, CyDs, are macrocyclic oligosugars most commonly composed of 6, 7, or 8 glucosidic units bearing the names  $\alpha$ -1,  $\beta$ -2 and  $\gamma$ -3 CyD, respectively [1–3]. Other, usually smaller, molecules (called guests) can enter their cavity forming inclusion complexes with these hosts.  $\alpha$ - and  $\beta$ -CyDs are believed to have been isolated for the first time at the end of nineteenth century [4] while their first complex seems to have been reported early in the twentieth century [5]. However, it took more than 50 years to establish and confirm the structure of CyDs. Today we take for granted the idea of inclusion complex formation by these macrocycles, but when it was suggested by Cramer in the 1940s the idea was not, to put it mildly, generally accepted. Cramer said later with considerable enjoyment [6]: “When I presented my results for the first time at a meeting in Lindau, Lake Konstanz, I met fierce opposition from some parts of the establishment. One of my older (and very important) colleagues even stated publicly and bluntly in the discussion that he would try to remove a young man with such crazy ideas from the academic scene. But there was also a good number of supporters, so I finally made it.”

According to Stoddart [6], “Cyclodextrins are all-purpose molecular containers for organic, inorganic, organometallic, and metalloorganic compounds that may be neutral, cationic, anionic, or even radical.” They are usually built of glucopyranoside units in the  ${}^4C_1$  conformation (Fig. 1.1). In most cases these host molecules have an average structure of a truncated cone with a cavity lined with H3 and H5 protons and lone pairs of glycosidic oxygen atoms lying in a plane thus endowing the cavity with hydrophobic character, while the bases formed by the primary and secondary OH groups bestow a hydrophilic character (Fig. 1.2). The great significance of CyDs both in research and applications lies in their ability to selectively form inclusion complexes with other molecules, ions, or even radicals. This phenomenon bears the name *molecular recognition* while the selectivity in the formation of complexes with enantiomeric species as guests is called *chiral recognition*. Complex formation changes the properties of both host and guest, allowing one to monitor the process by several experimental techniques. On the basis of X-ray



1

2



3






Fig. 1.1. Notation of conformations of the glucopyranoside ring.




**Fig. 1.2.** Schematic view of the glucopyranoside ring with the atom numbering and native CyD sizes and the average orientation of the most important atoms and OH groups.

measurements, native CyDs 1–3 have for years been considered to possess a rigid, truncated-cone structure [7–9]. This view is inconsistent both with the CyDs' ability to selectively complex guests of various shapes and with several experimental and theoretical findings, discussed later in this chapter. These data reveal the amazing flexibility of the CyD macrocycles. The implications of the non-rigidity of CyDs for their complexing ability, and its influence on the results obtained using different experimental techniques, are also presented there. One of the most striking examples of this kind is provided by the different mode of entrance of the guest in the complex of nitrophenol **4** with permethylated  $\alpha$ -CD **5** ( $R_1 = R_2 = R_3 = \text{Me}$ ) [10] in the solid state and in solution shown in Fig. 1.3.

The ability to predict recognition by CyDs would be of great practical value, especially for drug manufacturers. Consequently, several models of chiral recognition by CyDs have been proposed in the literature, neglecting the complexity of the complexation process involving very small energy differences between the complexes with enantiomeric species. The models critically reviewed later in this chapter are mostly based on very few experimental data and some of them contradict




**Fig. 1.3.** A schematic view of the entrance of nitrophenol guest **4** in the cavity of **5** ( $R_1 = R_2 = R_3 = \text{Me}$ ) in solution (a) and in the solid state (b).




5

the basic properties of three-dimensional space. For instance, the most often used 3-point Dalglish model [11] is incompatible with the basic requirement of three-dimensional space that at least four points (not lying in a plane) are necessary for an object to be chiral [12, 13].


Numerous CyD derivatives have been synthesized with the aim of improving their complexing properties and to make them suitable for various applications, in particular to increase the bioavailability of a drug complexed with a particular CyD derivative. By appropriate choice of host and guest one can achieve a very high selectivity. For instance, **6** is complexed much more strongly by the dimeric host **7** than is **8** [14]. Numerous CyD derivatives mono- or polysubstituted in positions 2, 3 and/or 6 by alkyl groups **5**, **9** as well as modifications of hydroxyl groups



6




7




8






9


to sulfopropyl, carboxymethyl, tosyl, aldehyde, silyl, and many other groups have been obtained [15, 16]. The reactivity of CyD and the plethora of exciting CyD structures developed, among other reasons, to enhance and modify their complexing ability will be shown in Chapter 2. Fascinating CyD structures include, among others, **10** [17], **11** [18], amphiphilic **5** ( $R_1 = R_2 = \text{OH}$ ,  $R_3 = \text{CH}_2\text{S}(\text{CH}_2)_3\text{C}_6\text{F}_{13}$  [19], capped **12** [20], peptide appended **13** [21] and 2:2 complex **14** formed by the CyD dimer with porphyrin and zinc ion [22]. On the other hand, obtaining dimers




10



11



12




n = 6, 7    R = Glycyl-L-Phenylalanyl

13

of isomeric naphthalenic acid, using an appropriately substituted  $\gamma$ -CyD template to exert control of the reaction's stereochemistry, shows a very elegant method making use of the encapsulation of two naphthyl-involving substituents on different glucopyranoside rings (Fig. 1.4) [23].

Exciting CyDs involving oligomers and polymers both covalently bound and self-assembled will be presented in Chapter 3 while their SPM observations and some




## 14

polymers having catenane or rotaxane structures (Fig. 1.5) will be discussed in Section 10.6 and Chapter 12.

CyD catalysis, discussed in Chapter 4, and the application of CyDs as enzyme models constitute a fascinating field. The influence of a specific CyD on the stability of the included molecule can have, like the two-headed god Janus, contrasting consequences. Mostly, luckily for pharmaceutical applications, complexation with CyD usually stabilizes the guest. (However, it can also catalyze its decomposition as is the case with aspirin **16** [24] preventing its application in the form of a CyD complex.) CyDs are known to catalyze numerous reactions. Notably, the catalytic activity is usually not high, but (a) it can reach high values in a few cases as evidenced by the ca. 1.3 million-fold acceleration of the acylation of  $\beta$ -CyD **2** by bound *p*-nitrophenyl ferrocinnamate **17** [25] and (b) CyDs may impose limitations on the reaction's regioselectivity. Chlorination of anisole in the absence or presence of  $\alpha$ -CyD illustrates this point [26] (Fig. 1.6) since it is known to produce only *para*-substituted isomers in the presence of  $\alpha$ -CyDs while both *meta*- and *para*-isomers are obtained in its absence [27]. It should be stressed that, although the catalytic activity of CyD can achieve high levels [25, 28], these oligosaccharides are much more effective in inducing stereo- or regioselectivity than in genuine catalytic action. Another example of regioselectivity induced by  $\gamma$ -CyD was presented earlier in Fig. 1.4.

An exciting field of considerable importance related to CyD catalysis is their use as enzyme models by testing the reactivity of appropriately substituted CyDs [29–32]. For instance, to mimic the cleavage of RNA followed by cyclization of phosphate ester with subsequent hydrolysis using imidazole groups of Histidine-12 and Histidine-119 of ribonuclease A, isomeric diimidazole-substituted at C6 positions  $\beta$ -CyDs **18–20** were synthesized [29] and checked for their influence on the



**Fig. 1.4.** Regioselectivity of the reaction of the included guest: schematic views of (a) disubstituted  $\gamma$ -CyDs, (b) the course of the reaction, (c) the product and yield of the reaction.

cleavage reaction. Interestingly, contrary to the classic mechanism proposed for this reaction, the close to linear arrangement of imidazole **21** groups in **20** did not lead to the most efficient catalysis, causing the abandonment of the mechanism commonly accepted in textbooks. The catalytic action of nuclease, ligase, phosphatase, and phosphorylase was also analyzed using more complicated CyD derivatives [33].










Fig. 1.5. Schematic catenane (a) and rotaxane (b) structures.







Fig. 1.6. Influence of  $\alpha$ -CyD on anisole chlorination.


As mentioned before, CyDs and their complexes elicit a vivid interest as systems that allowing us to study the factors that drive the selective complexation known as molecular recognition. Particularly great interest is focused on the differentiation between enantiomers of guest molecules by their complexation with CyDs, i.e. on chiral recognition. This will be of great importance in future CyD applications, in particular in the pharmaceutical industry, since most drugs and the active sites in which they operate are chiral. As a consequence (remember a small child trying to put his foot into the other shoe?), enantiomers of several drugs have been found to exhibit different pharmacological activities [34]. The differences may go so far as to result in harming instead of healing. (The old thalidomide **22** tragedy, when pregnant women taking this drug in the racemate form later gave birth to babies with crippled extremities, was sometimes interpreted in terms of the teratogenic activity of its second enantiomer [35]. However, the recent revival of interest in thalidomide drug for various illnesses [36–38] should be acknowledged.

Chromatography is one of the most important methods for direct studies of molecular and chiral recognition by CyDs. Today it has split into several branches, e.g. gas chromatography, GC, high-performance liquid chromatography, HPLC, and capillary electrophoresis and other electromigration techniques, that enable us not only to detect the recognition but also to estimate the complex stoichiometry and formation constant and, consequently, the enthalpies and entropies of complex for-




**22**

mation. The amazing sensitivity of CyDs to the shapes of guest molecules or ions may be illustrated by the big difference among retention times of the complex of 1,8-dimethylnaphthalene **23h** with **2** on the one hand and those of other isomers **23a–23g** on the other, determined by gas–liquid chromatography (discussed in Chapter 4 in more detail) [39]. Such a big difference is most probably caused by a difference in the stoichiometry of these complexes. Namely, the latter complexes are of 1:1 stoichiometry while in the former one guest molecule is mostly embedded in a capsule formed by two host CyDs [40]. The striking change in elution order with temperature rise indicates the importance of the entropy factor and of CyD flexibility on the complex stability [41]. Similarly, the dependence of the elution order of the enantiomers of phenothiazines **24** complexed with  $\gamma$ -CyD **3** on the buffer used shows the complexity of the complexation process [42]. Molecular and chiral recognition by CyDs, as studied mainly by HPLC and GC, will be presented in Chapter 5 together with the application of this method for studying complex stoichiometries and stability constants while a wide range of chromatographic methods used for enantioseparations will be discussed in Chapter 6.



In most cases, CyD structures are elucidated on the basis of X-ray studies which will be presented in Chapter 7 together with the results of a few, but very interesting, neutron diffraction investigations. They include systems of  $O2H \rightarrow O3$  and  $O3H \rightarrow O2$  hydrogen bonds in  $\beta$ -CyD **2** rapidly interchanging at room temperature (Fig. 1.7) [43]. Freezing the process and accurately determining the positions of hydrogen atoms using neutron diffraction [9] allowed the determination of the







**Fig. 1.7.** Rapidly interconverting circular systems of hydrogen bonds in  $\beta$ -CyD 2 (flip-flop, HO2  $\rightarrow$  O3 simultaneously and reversibly changing to O3H  $\rightarrow$  O2).

circular systems of the bonds shown in the figure. The mechanism of simultaneous change of directions of hydrogen bonds in 2 is called “flip-flop”. Other fascinating examples of X-ray determined CyD structures are provided among others by [3] catenane-type CyDs 25 [44], the sixteenfold deprotonated  $\gamma$ -CyD dimer with Pb ions 26 [45], and the complex of an alkali ion buried inside a capsule formed by two crown ethers, in turn inserted in another capsule built of two  $\gamma$ -CyD molecules, the whole system resembling a Russian doll [46, 47].


The forces driving complexation by CyDs cannot be understood without a knowledge of the energy differences and barriers involved in the complexation. The calorimetric measurements, involving isothermal titration calorimetry and differential scanning calorimetry are discussed in Chapter 8. They give the most accurate thermodynamic data characterizing the complexes. In particular, these data provide further examples of the amazing enthalpy–entropy compensation that is not limited to CyD complexes [48]. Exciting studies of isotope effects on complex formation are also discussed there.

X-ray and neutron diffraction studies yield precise information on the CyD structure in the solid state. On the other hand, in addition to the information on the structure and dynamics of complexes in the solid state, NMR spectra allow elucidation of the structure in solution, which is of particular importance since most CyD applications take place in this state. (Even if we take a drug as a CyD complex in the form of a pill it dissolves in the stomach before acting.) NMR studies can give not only unequivocal proof of the complex formation in form of, usually small, chemical shifts but also, by studying the NOE effect [49], they can show how the organic guest molecule enters the host cavity in the solid state and in solution. The spectra are also sensitive to the dynamics of the complex and so they provide



**25a****25b****26****27**

information on the complex's nonrigidity showing the host and/or guest movement even in the solid state where, owing to positional and time averaging, X-ray results point to a single rigid structure. This is the case for the complex of benzyl alcohol **27** with **2** for which  $^2\text{H}$  NMR spectra indicate a rapid flip of the aromatic ring around the C1C4 axis [50]. In addition to information on the complex's




stoichiometry and stability constants, in favorable cases the study of relaxation rates in  $^1\text{H}$  NMR spectra can show the orientation of the guest in the host cavity in solution, which no other technique can give, as shown for the complexes of camphor enantiomers **28** with **1** [51]. The kind of information on CyDs and their complexes that can be provided by NMR studies is discussed in Chapter 9 while Chapter 10 is devoted to the application of other physicochemical methods (UV, circular dichroism, mass spectroscopy, electrochemistry, AFM and STM, etc.) to the elucidation of the structure of CyDs and their complexes. Some of these methods are usually less sensitive to complex formation involving CyDs, but the effect can be considerable in specific cases and is of importance for applications in sensors and other devices. Although CyDs themselves do not have electroactive groups, electrochemical studies of their complexes form the basis of their future applications. Dendrimers with electroactive end groups like **29** [52] forming multiple CyD complexes are, probably, one of the most interesting examples in this area. Of course, mass spectra are most frequently used to prove the synthesis of a CyD derivative but, as shown in this chapter, they can be a source of valuable information on CyD complexes. New, rapidly developing AFM and STM techniques allowing the study of CyD aggregates on surfaces will also be presented there. They provide information on a single molecular aggregate or superstructures formed by


them. In particular, rotaxane-type structures **15b** (discussed in detail in Chapter 12) can be observed with atomic resolution by the latter method.

The possibility of predicting the molecular and chiral recognition ability of CyDs would be of great value, in particular for the pharmaceutical industry. The need for reliable theoretical treatment of CyD complexes is also reflected in several chapters in this book. Numerous studies applying quantum mechanics [53], molecular mechanics [54], and molecular dynamics [55] have been published by researchers fascinated by beautiful computer models and the ease of carrying out the calculations. The complexity of the complexation process and its consequences for the nonrigidity of CyDs, as well as the limited accuracy of the calculations, are neglected in most of these studies. Modeling of CyDs and their complexes and the dependence of the results of calculations on the assumed model and its parameterization are critically reviewed in Chapter 11.

The exciting catenanes, like **25** [44], and rotaxane molecular necklace **30** of 1:*n* stoichiometry incorporating  $n = 20\text{--}22$   $\alpha$ -CyD macrocycles [56], respectively, falling into the realm of topological chemistry [57] will be shown in Chapter 12. These systems, also discussed in Chapters 10.6 and 16, form the basis of exciting applications. Large CyDs such as the 12-membered **31**, which differ dramatically from native CyDs in properties and, most probably, in complexing ability, will be discussed in Chapter 13. In particular, contrary to the structure of **1–3**, large CyDs do not have truncated-cone average structures with glycosidic oxygen atoms lying approximately in a single plane, but some of them are known to be twisted allowing for formation of hydrogen bonds between OH groups of distant glycosidic units [58].



Chiral recognition by CyDs is of primary importance for the pharmaceutical industry since the second enantiomer of a drug, usually present as 50% impurity as the result of chemical synthesis, can be harmful. Therefore, an effective preparative separation of enantiomers is one of the important goals of applied CyD research since at present it has not reached the industrial scale. Today the main CyD application in the pharmaceutical industry is their use as drug carriers, since CyD containers in most cases stabilize and solubilize the included drugs (see, how-



### 31

ever, the aspirin case mentioned above). Moreover, the slow release of a drug from the complex results in its higher and more uniform content in the organism, allowing less frequent administration of the drug. Interestingly, CyD applications in drug delivery were considerably delayed by the erroneous determination of their toxicity at an early stage of development [59]. Today we know that they are not harmful in most cases by oral, parenteral, nasal, or skin administration [60]. CyD applications in the form of inclusion complexes in the pharmaceutical industry in general are presented in Chapter 14 with a detailed discussion of the ways in which various CyD types (hydrophilic, hydrophobic, or ionizable ones) affect the bioavailability of drugs by influencing their solubility and the rate of release from the CyD complex. A small section on site-specific drug delivery is also included. The even better therapeutic effect of drugs in the form of emulsions, micro-particles, nanoparticles, and higher aggregates is given in Chapter 15.

CyD applications are by no means restricted to the pharmaceutical industry. Several examples, mainly prospective ones, are scattered throughout this book. CyDs are used to remove unpleasant tastes, odors, or other undesirable components in the food industry, in agrochemistry, cosmetics, dyeing, cleaning, and in many other areas. To name just a few examples of numerous CyDs applications: grapefruit juice loses its unpleasant taste when its bitter component naringine is removed by complexation with  $\beta$ -CyD [61]; similarly, removal of phenylalanine and tyrosine

makes the food harmless for those suffering from phenylketonuria [62]. Garlic retains all its health benefits but lacks its annoying odor when applied in the form of a CyD complex. Similarly, other flavor components (such as apple, citrus fruits, and plums) and spices (cinnamon, ginger, horseradish, menthol, etc.) are marketed as  $\beta$ -CyD complexes characterized by high stability exhibited towards heating during industrial food processing [63–66]. The problems encountered by the use of CyDs for stabilization of mixtures has to be mentioned here since the components involved can be released at different rates, changing the taste or smell of the product. The creation and higher stability of foams used in both the food and cosmetic industries is also favored by the complexation. Interestingly, CyD applications should not only improve the properties of a marketed product but, consistently with modern trends, they are also aiming at creating new needs by producing new types of products with unheard of properties. A long-lifetime fragrance-releasing room-decorating paint is one example of such applications [67]. A few examples of numerous CyD applications in industries other than pharmaceuticals are briefly presented in Chapter 16, which concludes with the presentation of their thought-challenging prospective applications in molecular devices and machines.

The cyclodextrin field is rapidly expanding. According to information from the Cyclolab website, 3.9 articles per day on CyDs were published on average in 1995. This figure increased to 4.4 in 2004 ([http://www.cyclolab.hu/literature\\_0.htm](http://www.cyclolab.hu/literature_0.htm)) and is still growing. The field is not only very large but also highly diversified. This is probably the reason why no general modern comprehensive monograph on CyDs has been available [1–3, 16, 68–71] for so long. We would like to bridge this gap, providing a survey of the whole exciting area with numerous references to help novices to enter this domain on the one hand and to give researchers in a specific field a broader insight into the whole area on the other. We apologize for not being able to mention, even briefly, numerous valuable works on CyDs but we hope to have given the picture of the whole CyDs field, illustrated by representative examples of the respective research and applications.

## 1.2 Cyclodextrin Properties

In the standard way, native CyDs 1–3 are obtained by enzymatic degradation of starch. First obtained in minute amounts and very expensive, in particular  $\alpha$ - and  $\gamma$ -CyD, they now cost less than \$10/kg, making their large-scale industrial use feasible [72]. IUPAC names of these macrocycles are cumbersome; 5,10,15,20,25,30,35-heptakis-(hydroxymethyl)-2,4,7,9,12,14,17,19,22,24,27,29,32,34-tetradeca-octa-cyclo[31.2.2.2.<sup>3,6</sup>.2<sup>8,11</sup>.2<sup>13,16</sup>.2<sup>18,21</sup>.2<sup>23,26</sup>.2<sup>28,31</sup>]nonatetracontane-36,37,38,39,40,41,42,43,44,45,46,47,48,49-tetradecol for  $\beta$ -CyD makes the use of trivial names necessary. Lichtenthaler and Immel proposed a general system of naming macrocyclic oligosaccharides [73] but it has not been generally accepted and used. The chemical, physical, and biological properties of CyDs, and in particular their toxicity by various type of administration, are summarized in Ref. [60] while their stability when


Table 1.1. Some properties of native CyDs [60]<sup>a</sup>

	$\alpha$ -CyD 1	$\beta$ -CyD 2	$2\gamma$ -CyD 3
Number of glucose units	6	7	8
Molecular weight	972	1134	1296
Approximate inner cavity diameter (pm)	500	620	800
Approximate outer diameter (pm)	1460	1540	1750
Approximate volume of cavity ( $10^6 \text{ pm}^3$ )	174	262	427
$[\alpha]_D$ at 25 °C	$150 \pm 0.5$	$162.5 \pm 0.5$	$177.4 \pm 0.5$
Solubility in water (room temp., g/100 mL)	14.5	1.85	23.2
Surface tension (MN/m)	71	71	71
Melting temperature range (°C)	255–260	255–265	240–245
Crystal water content (wt.%) <sup>*</sup>	10.2	13–15	8–18
Water molecules in cavity	6	11	17

<sup>a</sup> Some data on these and larger CyDs are also given in Tables 9.1 and 13.1.

treated by various enzymes is presented in Ref. [74]. Some of their properties are given in Table 1.1.

As mentioned earlier, native CyDs are usually obtained by biochemical processes [72]. However, a 21-step synthesis of **1** with 0.3% total yield [75] and that of **3** [76] with 0.02% yield are worth mentioning. The macrocycle built of only five glucopyranose rings **32** [77] (thought to be too strained to exist on the basis of model calculations [78]) and probably thousands of CyD derivatives have been synthesized [15, 16]. Several exciting CyDs have been presented earlier. Here CyDs having glucopyranoside ring(s) in  ${}^1C_4$  [79–81] or *skew* conformations [82], those incorporating other than glucopyranoside units [83] and large CyDs having from 9 to

**32**

more than 100 monosaccharide rings (discussed in Chapter 13) should be mentioned [58, 84, 85]. Interestingly, three unusual CyD derivatives have been found in nature [86].

As discussed in Chapter 7, native CyDs can form complexes with differing amounts of water. CyDs are seldom really empty. Even if they do not contain another guest there is usually at least one solvent molecule in their cavity. Almost all their applications involve inclusion complex formation when one or more molecules are at least partly immersed in the CyD cavity. The complexes can be obtained both in solution (sometimes requiring heating or the use of a cosolvent) or in the solid state, e.g. by cogrinding or milling [87]. In solution, they exist in a rapidly exchanging equilibrium of the free CyD host and guest. As mentioned before, CyD complexes of different stoichiometry are known. In addition to 1:1 complexes like **4@5** ( $R_1 = R_2 = R_3 = \text{Me}$ ) [10], the 1:2 ones like those of camphor enantiomers **28** embedded in a capsule formed by two  $\alpha$ -CD **1** [51] and that of  $C_{60}$  buried in two  $\gamma$ -CyD molecules **3** in a similar way [88], 2:2 complex **14** [22] or even rotaxane molecular necklace **29** of 1: $n$  stoichiometry involving  $n = 20$ – $22$   $\alpha$ -CyD macrocycles [56] are known. In spite of the years that have passed from the publication of the Szejtli review [89], his statement “The ‘driving force’ of complexation, despite the many papers dedicated to this problem, is not yet fully understood.” is still valid. The complexation process in a, mostly water, solvent is considered to involve a release of water molecule(s) from the relatively hydrophobic CyDs’ cavity, removal of the polar hydration shell of the apolar guest molecule, entry of the guest into the empty CyD cavity where it is stabilized mainly by weak but numerous van der Waals attractive interactions, restoration of the structure of water around the exposed part of the guest, and integration of this with the hydration shell of host macrocycle. Thus, a change in both enthalpic and entropic contributions occurs in complex formation that depends on the host- and guest-induced fit [90], the solvent used, and numerous other factors. On the other hand, in the solid state the magnitude of the crystal forces is comparable with the forces keeping the complex together. Thus, as exemplified by **4@5** ( $R_1 = R_2 = R_3 = \text{Me}$ ) [10], they can influence the complex’s structure and dynamics which is considerable even in the solid state. To summarize: CyD complexes are very difficult to study since (1) for poorly soluble species the complexation process can be much more effective for impurities (present in minute amount in the solution) than for the guest under investigation; (2) the process depends heavily on the experimental conditions (pH, cosolvent, temperature, etc.); (3) the complexes can involve species of different stoichiometries, e.g. dimethylnaphthalenes **23a–g** and **23h** [40], or an additional solvent molecule can enter the cavity as a second guest resulting in ternary complex formation [91, 92] (interestingly, complexes of even higher stoichiometry, involving two CyDs, one pyrene, and two cyclohexanol guest molecules [92] are known); (4) the experimental results for a CyD complex can depend on the technique used since the CyD complexes are held together by weak forces (one example of this kind was shown in Fig. 1.3 [10]); (5) Reliable theoretical studies for CyD complexes are extremely difficult to obtain since, as discussed in detail in Chapter 11, these large systems are characterized by energy surfaces exhibiting numerous very shal-

low local minima separated by low energy barriers [93, 94]. The size of the system and its  $n$ -fold degeneracy ( $n = 6, 7, 8$  for  $\alpha$ -,  $\beta$ -, or  $\gamma$ -CyD, respectively) also make it difficult to compare X-ray geometrical parameters for, for example, complexes with different guests. As discussed in detail in Section 1.4, one can either attempt to analyze the values of all internal parameters (e.g. of 126 bond lengths in  $\alpha$ -CyD, etc.) in the series, which gives a very nontransparent picture, or compare the values of, for instance, the C2O2 bond length averaged over all saccharide rings, losing a lot of information by such an approach.

### 1.3

#### Cyclodextrin Nonrigidity [94, 95]

On the basis of X-ray studies [7–9], for tens of years CyDs structure was thought to resemble a rigid truncated cone of the high  $C_6$ ,  $C_7$ , or  $C_8$  symmetry for  $\alpha$ -,  $\beta$ -, or  $\gamma$ -CyD, respectively, with a planar ring of glycosidic oxygen atoms [8, 96, 97]. Numerous experimental and theoretical data are incompatible with the concept of rigid CyDs. First of all, a general analysis of these systems shows that there is no physical reason for the rigidity, since the macrocycles are built of relatively rigid glucopyranose rings connected by ether C–O bonds, characterized by a low barrier to internal rotation of ca. 1 kcal mol<sup>-1</sup> [98]. This reasoning was supported by model molecular mechanics calculations on  $\alpha$ -CyD [93] showing that (a) the usually depicted structure with planar rings formed by glycosidic oxygen atoms does not correspond to the energy minimum and (b) the energy hypersurface exhibits several energy minima separated by low barriers. With regard to CyD complexes, they are held together by weak intermolecular interactions which somewhat limit the macrocycle's mobility but cannot endow the macrocycle with considerable rigidity.

It should be emphasized that a rigid structure for CyDs is also incompatible with the ease of formation of inclusion complexes of various shapes, since the latter implies an effective fitting of the host and guest to each other [90]. Most experimental proofs of the nonrigidity of CyDs come from NMR studies not only in solution but even in the solid state. If CyDs were not flexible then the spectra of complexes with aromatic guests in solution should exhibit several signals for, e.g. H3 CyD protons on different glucopyranose rings pointing into the cavity (Fig. 1.2). This is not the case [99]. Moreover, NMR studies in the solid state show that the rings included in the CyD cavity can exhibit a rapid flip around the C1C4 axis. One example is provided by **27@2** for which <sup>2</sup>H NMR spectra are incompatible with the rigid structure [50]; similar evidence has been obtained on the basis of <sup>13</sup>C NMR spectra [100, 101]. The rapid inversion of *cis*-decalin **33** in the complex with **2** at room temperature frozen at 233 K, observed in both <sup>1</sup>H and <sup>13</sup>C NMR spectra [102, 103], is also incompatible with CyD rigidity.

The very fast internal movement of native CyDs and of most of their derivatives, leading to the observation of averaged structures by most experimental techniques, is frequently overlooked. In addition to the temperature-dependent process of self-inclusion of substituent(s) [104–107], we were able to find only two studies of substituted CyDs in which movement of the macrocycles was at least partly frozen [104, 108]. Some other experimental results proving CyD flexibility using NMR